










Photos by Peter LaTourrette and PRB

Predicting Effects of Habitat Change on Avian Communities
Diana Stralberg, Nils Warnock, Dave Shuford, Sam Valdez

Outline

- Salton Sea habitat conversion model
 - Modeling objectives and approach
 - Focal species
 - Major habitats and model variables
- San Francisco Bay habitat conversion model
 - Background
 - Sample results from Phase I

Salton Sea Habitat Conversion Model



PRBO Modeling Objectives

- Evaluate and compare the effects of specific habitat configuration alternatives on waterbirds.
- Identify key habitat features that may be manipulated to enhance waterbird numbers.
- Identify important landscape characteristics that affect waterbird habitat value.
- Identify optimal habitat configurations.

PRBO Modeling Approach

- Empirical models
- Static, habitat-based
- Assume habitat is limiting
- Limited to habitat / landscape characteristics that are easily quantified
- Predictions represent population indices for focal species and habitat-guilds

Modeling Steps

- Develop statistical models based on 1999 bird data, field-collected habitat data, and GIS-based habitat / landscape data
- Provide information on important habitat and landscape characteristics
- Evaluate specific habitat alternatives
- Provide predictions of bird numbers (population indices) under each alternative (by habitat / guild and individual focal species)

Waterbird Focal Species by Habitat-Guild

Species	List	Open Water		Shoreline			Snags /	Islands /	Freshwater	Saline	
		Fish	Inverts	Fish	Inverts	Deltas	Trees	Dikes	Marsh	Ponds	Agriculture
Eared Grebe	Α		Х							Х	
Brown Pelican	FE, SE	Х						Х			
American White Pelican	BSSC, A	Х		Х		Х		X			
Double-crested Cormorant	BSSC, A	Х				Х	Х	X	X		
Snowy Egret	WP			Х			Х		X		X
Cattle Egret	Α						Х		X		X
Least Bittern	BSSC								X		
White-faced Ibis	BSSC, A					Х	Х		X		X
Wood Stork	BSSC, WP			X	X	Χ	X		X		
Fulvous Whistling Duck	BSSC								X		
Ruddy Duck	Α		Х								
Redhead	BSSC		X		Х	Х			X	X	
Black Rail	ST, BCC								X		
Yuma Clapper Rail	FE								X		
Greater Sandhill Crane	ST										X
Snowy Plover	BSSC, SP, BCC				X						
Mountain Plover	BSSC, SP, BCC										Х
Black-necked Stilt	Α				Х	Х		X	X	Х	X
American Avocet	Α				Х	Х		X	X	X	X
Long-billed Curlew	BSSC, SP									X	X
Whimbrel	SP, BCC									Х	X
Marbled Godwit	SP, BCC				Х	Х				Х	
Western Sandpiper	SP				Х	Х				X	
Dunlin	SP				Х	Х				X	
dowitcher spp.	Α				Х	Х				Х	
Ring-billed Gull	Α			Х	Х	Х		X			X
California Gull	Α			Х	Х	Х		X			
Gull-billed Tern	BSSC, WP, BCC			Х	Х	Х		X			X
Black Tern	BSSC	Х		Х	Х	Х		X		Х	X
Black Skimmer	BSSC, BCC, WP	Х		X		Х		Х			
Burrowing Owl	BSSC, BCC										Х
BSSC = Calif. Dept. Fish and Game Bird Species of Special Concern;						SP = US Shorebird Conservation Plan species with a 4-5 priority score					
SE = state endangered; ST = State threatened					A = >10,000 birds counted on single survey (Shuford et al. 2002)						
FE = Federally Endangered;						BCC = USFWS Birds of Conservation Concern - BCR 33					
WP = National Waterbird C		ecies con	sidered High	nly Imperi	led or of Hi	gh Concern	1				

Shoreline / Shallow Water Bird Data


 Comprehensive shoreline surveys (Jan, Apr, Aug, Nov 1999)



Shoreline / Shallow Water Habitat Variables

- Shallow water characteristics (within 500 m of shoreline)
 - min, max, mean elevation (depth)
 - shallow proportion (< 1 ft)
 - mean slope
- Shoreline characteristics
 - shoreline complexity
 - proportion of wetlands within 0.5 1 5 km
 - proportion of agricultural land uses within 0.51 5 km
 - distance to nearest primary (deltas) and secondary (creeks) river mouth

Open Water Bird Data



Open Water Habitat Variables


- Min, max, mean elevation (depth)
- Substrate characteristics (?)
- Distance to nearest primary (deltas) and secondary (creeks) river mouth

Managed Marsh Bird Data

Comprehensive wildlife area /

refuge / duck club surveys


Clapper Rail surveys



Managed Marsh Habitat Variables

- Open water proportion (?)
- Vegetation height (?)
- Marsh size
- Proportion of agricultural land uses within
 0.5 1 5 km
- Distance to nearest primary (deltas) and secondary (creeks) river mouth

Agricultural Bird Data




Agricultural Habitat Variables

- Crop type
- Crop height
- Crop status (mowed, tilled, growing)
- Moisture (dry, partial, or full flooding)
- Proportion of general crop types along transect (leaf, bulb, alfalfa, grass, bare, unsuitable)

Managed Pond Bird Data


San Francisco Bay Salt Pond Surveys




Managed Pond Habitat Variables

- Mean depth
- Shallow proportion (< 15 cm)
- Deep proportion (> 1 m)
- Salinity (ppt)
- Pond size
- Surrounding land use within 1 km

South San Francisco Bay Habitat Conversion Model




Historic Habitat Change: Diking, Filling, Dredging




Current Habitat Change: Salt Pond Restoration

- 5,471 ha of salt ponds purchased for restoration.
- Restoration and management decisions will involve trade-offs for birds.



Restoration Trade-offs




Salt Ponds



Tidal Marshes

Bird Use of Tidal and Managed Wetlands




Site Characteristics



Landscape Characteristics

- Distance to open bay, development
- Surrounding landscape composition



Site-level Restoration Scenarios*

1: Salt ponds retained at current salinities (no change)

All salt ponds restored to tidal marsh:

2: minimum large channels and ponded areas



3: mean large channels and ponded areas




4: maximum large channels and ponded areas




^{*} Landscape conditions incorporated into models.

Landscape-level Restoration Scenarios






Scenario Comparisons, Small Shorebirds





Scenario Comparisons, Diving Ducks



San Francisco Bay Phase I Conclusions

- Loss of managed ponds may cause substantial reduction in waterbird numbers, especially small shorebirds and diving ducks.
- Potential to reduce and/or avoid waterbird losses through design and management of individual restoration sites.
- Critical for waterbirds to retain some characteristics of managed ponds in a habitat mosaic (more important than tidal marsh design and management).

San Francisco Bay Phase II Activities

- Identification of focal species and key seasons
- Incorporation of salt pond bathymetry data
- Explicit consideration of marsh microhabitats (pond/pan, channel, vegetation)
- Incorporation of intertidal mudflats, open bay
- Approximation of carrying capacities by guild
- Use of information theory and model selection
- Evaluation of realistic restored marshes
- Evaluation of optimally managed ponds

Acknowledgements

Funders:

California Coastal Conservancy
Gabilan Foundation
Mary A. Crocker Trust
Rintels Charitable Trust
Tides Foundation

Partners:

San Francisco Bay Bird Observator
U.S. Geological Survey
U.S. Fish & Wildlife Service
(Don Edwards San Francisco Bay NWR)

